

**Test 2 for Calculus III for CS Majors, Math 2506 J1-J2, October 23, 2007**

**Name:**

This test is to be taken without calculators and notes of any sort. The allowed time is 50 minutes. Provide exact answers; not decimal approximations! For example, if you mean  $\sqrt{2}$  do not write 1.414... State your work clearly, otherwise credit cannot be given.

**Problem 1:** (10 points) a) Diagonalize the matrix  $A = \begin{bmatrix} 1 & 4 \\ 1 & -2 \end{bmatrix}$ , i.e., find an invertible matrix  $S$  and a diagonal matrix  $D$  so that  $A = SDS^{-1}$ .

b) (10 points) Find a Schur factorization for the same matrix  $A$ , i.e., find an orthogonal matrix  $Q$  and an upper triangular matrix  $T$  so that  $A = QTQ^T$ .

**Problem 2:** Consider the matrix

$$A = \begin{bmatrix} 6 & 0 & 4 \\ 0 & 3 & 1 \\ 4 & 1 & 0 \end{bmatrix}$$

a) (5 points) Compute  $\text{Off}(A)$ .

b) (10 points) Calculate the Givens matrix  $G$  for the first step in the Jacobi iteration for diagonalizing  $A$  by picking the  $2 \times 2$  submatrix with the largest off diagonal elements. (You do not have to calculate  $G^T A G$ .)

c) (10 points) Calculate  $\text{Off}(G^T A G)$ .

**Problem 3:** a) (15 points) Calculate the singular value decomposition of the matrix

$$A = \frac{1}{3\sqrt{2}} \begin{bmatrix} 6 & 2 \\ 3 & 5 \\ 0 & 4 \end{bmatrix}$$

b) (10 points) Find the best rank one approximation.

**Problem 4:** Assume that

$$V = \frac{1}{3} \begin{bmatrix} 1 & 2 \\ -2 & 2 \\ 2 & 1 \end{bmatrix}, D = \begin{bmatrix} 3 & 0 \\ 0 & 2 \end{bmatrix}, U = \frac{1}{\sqrt{5}} \begin{bmatrix} 1 & -2 \\ 2 & 1 \end{bmatrix}$$

form the singular decomposition of a matrix  $B$ .

a) (10 points) Calculate the generalized inverse  $B^+$

b) (5 points) Find the least square solution of the problem  $Bx = b$  where

$$b = \begin{bmatrix} -2 \\ 1 \\ 2 \end{bmatrix}$$

**Problem 5:** (15 points) Find the maximum and minimum values of the function  $f(x, y) = x^3 + y^3$  in the region consisting of all points  $(x, y)$  that satisfy  $x^2 + y^2 - xy \leq 1$ . Find all the points where the maximum is taken and all the points where the minimum is taken. (Hint for the calculation: The identity  $x^3 - y^3 = (x - y)(x^2 + y^2 + xy)$  might be useful.)

**Extra Credit:** (10 points) Consider the matrix

$$A = \begin{bmatrix} 0 & 0.1 & -0.1 \\ 0.2 & 1 & 0.1 \\ -0.2 & 0.1 & 2 \end{bmatrix}$$

Find upper and lower bounds on the eigenvalues using the ‘Small Gershgorin Disk Theorem’.