Test IV for Calculus II, Math 1502, November 10, 2009

**Print Name:** 

Section:

Name of TA:

This test is to be taken without calculators and notes of any sorts. The allowed time is 50 minutes. Provide exact answers; not decimal approximations! For example, if you mean  $\sqrt{2}$  do not write 1.414.... Show your work, otherwise credit cannot be given.

Print your name (not your signature), your section number as well as the name of your TA on EVERY PAGE of this test. This is very important.

# Section:

# Name of TA:

**I:** (10 points each) For the two systems of equations below, decide whether solutions exist or not. If they do exist calculate all of them. a)

$$x + 2y = 3$$
  
$$2x + 3y - z = 4$$
  
$$2x + y - 3z = 1$$

b)

$$x + 2y = 3$$
  

$$2x + 3y - z = 4$$
  

$$2x + y - 3z = 0$$

#### Section:

# Name of TA:

**II:** Consider the matrix  $A = [\vec{v}_1, \vec{v}_2, \vec{v}_3]$  where

$$\vec{v}_1 = \begin{bmatrix} 1\\2\\1 \end{bmatrix}$$
,  $\vec{v}_2 = \begin{bmatrix} -3\\-4\\-1 \end{bmatrix}$ ,  $\vec{v}_3 = \begin{bmatrix} 2\\3\\1 \end{bmatrix}$ .

a) (5 points) Give a spanning set for Img(A).

b) (5 points) Give a basis for Img(A).

c) (10 points) Find a basis for Ker(A).

Section:

Name of TA:

**III:** True or false: (5 points each)

- a) For a symmetric matrix A, i.e.,  $A = A^T$ ,  $\text{Ker}(A)^{\perp} = \text{Img}(A)$ .
- b) An  $n \times m$  matrix whose rank is r has always n r non-pivotal columns.
- c) If the rank of a matrix A is r, so is the rank of the matrix  $A^T$ .
- d) Four vectors in  $\mathcal{R}^4$  are linearly independent if any three of them are.
- e) An  $n \times n$  matrix A is invertible if and only if  $\text{Ker}(A) = \{\vec{0}\}.$

**IV:** a) (15 points) Using the normal equations, solve the least square problem for  $A\vec{x} = \vec{b}$  where

$$A = \begin{bmatrix} 1 & 3\\ 2 & 1\\ 2 & -1 \end{bmatrix} , \quad \vec{b} = \begin{bmatrix} 2\\ 1\\ 1 \end{bmatrix}$$

b) (5 points) Find the vector in Img(A) that is closest to the vector  $\vec{b}$ .

#### Section:

# Name of TA:

**V:** A matrix A has the QR factorization with

$$Q = \frac{1}{3} \begin{bmatrix} 2 & -1 \\ 2 & 2 \\ 1 & -2 \end{bmatrix} , \text{ and } R = \begin{bmatrix} 3 & 1 & -1 \\ 0 & -1 & 1 \end{bmatrix}$$

i) (10 points) Find the orthogonal projection onto Img(A).

ii) (5 points) Find the orthogonal projection onto  $\text{Ker}(A^T)$ .

iii) (Extra credit: 15 points) Find the least square solutions for  $A\vec{x} = \vec{b}$ where  $\vec{b} = \begin{bmatrix} 4\\1\\-1 \end{bmatrix}$ .